

# AGRARES STICK & FAT







# **CHARACTERISTICS:**

AGRARES STICK it's a new formulation, rich in phosphorus and nitrogen, which incorporates molybdenum and boron in his composition, which applied by leaf favors the flowering and filling/fattening of the fruits. In addition, it reactivates the processes at cellular level that they favor the stick of the fruits in the branch of the tree, avoiding the falling.

# **DOSE:**

# **Product for LEAF application**

CROP	LEAF DOSE	MOMENT OF APPLI- CATION
Citric and fruit trees	100 cc/100L of water	To apply at beginning of flowering repiting every 10 days.
Horticulture (tomatoe, pep- per)	100 cc/100L of water	
Cucumber, me- lon, watermelon	100 cc/100L of water	
Other crops	100 cc/100L of water	



AGRARES iberia SL B-64.816.895

ALMACENES: Ctra. Logroño-Zaragoza Km 70, Nave6 26540 Alfaro, La Rioja ESPAÑA CONTACTO: agrares@agrares.com http://www.agrares.com Tel: (+34) 618 518 702 FAX: 940462034



Total Nitrogen6 % w/w (7,2% w/v)		
Ureic Nitrogen4,5 % w/w (5,4% w/v)		
Ammonia Nitrogen		
Anhídride phosphoric(P2O5)soluble in water.12 % w/w(14,4% w/v		
Microelements:		
Boron(B) soluble in water1% w/w (1,2% w/v)		
Molybdenum (Mo) soluble in water0,2% w/w (0,24 % w/v)		
Density1,2 gr/cc		
PH4		

# **COMPATIBILITIES:**

AGRARES STICK &FAT is compatible with most of the fertilizers NPK of normal use, not to mix with acid products (sulphuric, phosphoric, nitric), with products of alkaline reaction, with products sulphurcalcic not with mineral oils.

# **WARNINGS**:

S 36/37: To use gloves of protection and appropriate apparel.

To support far from food, drinks and piensos.

Be kept out of the scope of the children.

In case of contact with the skin, wash immediately and abundant with water.

To avoid the contact with the humid parts of the body, specially the eyes.

In case of contact with the eyes, wash abundant with water and go immediately to a doctor.

In case of accident or discomfort, come immediately to the doctor (if it is possible, show him the etiquette) not to store in another packing different from the original one.

Not to apply in hours of maximum sunstroke.

In case of lacking experience in mixtures consult to our Technical Department.

Be in use only in case of recognized need. Not to exceed the suitable doses.

# Bottles of:

1L, 5L, 20L, 200L y 1000L